

Reactive nitrogen species in cellular signaling

Levi Adams, Maria C Franco and Alvaro G Estevez

Burnett School of Biomedical Sciences, College of Medicine, University of Central Florida, Orlando, FL 32827, USA
Corresponding author: Alvaro G. Estevez. Email: alvaro.estevez@ucf.edu

Abstract

The transduction of cellular signals occurs through the modification of target molecules. Most of these modifications are transitory, thus the signal transduction pathways can be tightly regulated. Reactive nitrogen species are a group of compounds with different properties and reactivity. Some reactive nitrogen species are highly reactive and their interaction with macromolecules can lead to permanent modifications, which suggested they were lacking the specificity needed to participate in cell signaling events. However, the perception of reactive nitrogen species as oxidizers of macromolecules leading to general oxidative damage has recently evolved. The concept of redox signaling is now well established for a number of reactive oxygen and nitrogen species. In this context, the post-translational modifications introduced by reactive nitrogen species can be very specific and are active participants in signal transduction pathways. This review addresses the role of these oxidative modifications in the regulation of cell signaling events.

Keywords: Nitric oxide, nitrosylation, nitration, cell signaling

Experimental Biology and Medicine 2015; 240: 711–717. DOI: 10.1177/1535370215581314

Introduction

Transitory addition or removal of a prosthetic group allows a signal to be passed by inducing a change in the target molecule that alters its activity. Phosphorylation is the most widely discussed form of signal transduction and kinase cascades are very well characterized. Recently, reactive nitrogen species have been gaining attention as signal transduction mediators rather than damaging oxidizers of macromolecules. Oxidative modifications are difficult to study because the variety of targets makes it difficult to distinguish between relevant modification and collateral damage. In addition, when added exogenously, the concentrations of oxidants necessary to see consistent results are often damaging to the cell causing the concept of oxidation to become synonymous with cellular damage and senescence. However, the concept of oxidative signaling is now well established in the literature for a number of reactive oxygen and nitrogen species. The strongly oxidative nitrogen-based molecules and radical products of their decomposition are seen as too reactive to have the specific functions required for signal transduction. In addition, the modifications may be permanent causing them to be excluded from studies of signal mechanisms.¹ However, nitrated proteins are removed by degradation² and possible other mechanism yet to be fully characterized.^{3–6} In addition, new technologies that allow the study of proteins with single residue modification in the absence of

undesirable oxidative modification or damage to other cell components provide new opportunities for the investigation of oxidative signaling mediated by tyrosine nitration.^{7–9}

Oxidative modifications are often unstable and prone to further oxidation or interactions with reducing agents. On the other hand, the conditions of analysis can produce oxidative modifications that were not there in the first place increasing the difficulty for the analysis of oxidative modification. However, the field of redox proteomics has developed a number of methodologies to deal with these limitations. Combinations of biotin-based techniques, alkylating agents, and indirect fluorescent labeling can be used to find and “trap” the redox modifications, which can be identified with improved antibodies or without gels by using liquid chromatography and mass spectrophotometry. The multidimensional approach of redox proteomics can help provide insight into the mechanisms of endogenous redox signaling, and help provide diagnostic tools as we move forward.^{10–13}

Nitric oxide, nitrosation, and nitrosylation

Post-translational modifications involving reactive nitrogen species share a common progenitor: nitric oxide (NO). Nitric oxide is produced from L-arginine by three main isoforms of nitric oxide synthase (NOS): epithelial NOS (eNOS), related to vasodilation and vascular regulation; neuronal NOS (nNOS), which is linked to intracellular

signaling; and inducible NOS (iNOS), which has a variety of situational functions. While nitric oxide production by nNOS and eNOS is tightly regulated by calcium by a calmodulin-dependent mechanism,^{14,15} iNOS is activated in response to various endotoxin or cytokine signals, which can lead to the rapid production of large fluxes of nitric oxide. iNOS expression is regulated by well characterized signal pathways including MAPK and JNK/STAT,^{16,17} suggesting that the inducible production of nitric oxide must be tightly controlled. Several disease states have been linked to the deregulation of nitric oxide production, indicating that aberrant production of nitric oxide and its products can have deleterious consequences for the cells.^{18–23} All products formed by nitric oxide reactions are collectively denominated reactive nitrogen species, which include a number of compounds with very different chemical properties and reactivity.

Nitric oxide is a very versatile molecule with multiple functions and mechanisms of action. Soon after the discovery of nitric oxide it became evident that it could have opposing effects. Nitric oxide was described as a diffusible radical that results in vasodilation and a key player in the circulatory system. The Nobel Prize winning work of Murad et al. identified nitric oxide as ligand of the soluble guanylyl cyclase, which stimulates the production of cGMP.^{14,15,24} Nitric oxide-dependent production of cGMP has wide variety of targets, and plays a role in the regulation of several functions in the nervous system.^{25–32} However, it soon became clear that not all activities of nitric oxide were mediated by production of cGMP. Oxidative products of nitric oxide were soon reported in macromolecules and a number of proteins.

Early on, nitrosylation of thiols in cysteine residues was accepted as a possible post-translational modification,³³ often linked to reactions with oxygen or glutathione.^{1,34} Several mechanisms of nitrosylation have been described, including oxidative S-nitrosation, trans-nitrosylation by small molecular weight nitric oxide carriers like S-nitrosoglutathione, and metalloprotein-catalyzed S-nitrosylation.^{35–37} Nitrosylation occurs in a functionally diverse group of proteins in diverse subcellular locations and in different conditions, regulating a variety of cellular processes.^{36–38}

The first clearly defined role of nitrosylation in the nervous system was identified 20 years ago with the discovery that nitric oxide regulation of the NMDA receptor was mediated by S-nitrosylation.^{39–41} Lipton et al. identified nitric oxide produced by nNOS as both a positive and negative regulator of the NMDA glutamate ionotropic receptor, depending on redox state. Direct S-nitrosation of the NMDA receptor suppresses its activity, but reaction with the spontaneously formed nitric oxide/superoxide product peroxynitrite activates the receptor.⁴⁰ Another early target of nitrosylation is p21ras, which plays a central role in this paradoxical activity. The nitric oxide-mediated S-nitrosylation status of the NMDA receptor helps to regulate p21ras³⁹ and it was suggested that Ras is one of the central steps in how redox signals transmit their messages.⁴¹ Recent work showed that introduction of GSNO as a nitric oxide donor transiently nitrosylates Ras and Src kinase at the plasma

membrane, and is reversed within 20–40 min. Src kinase induces the diacylglycerol pathway, causing translocation of RasGRP1 to the Golgi apparatus, activating Golgi-associated Ras. In contrast, eNOS-generated nitric oxide induced by EGF only activates Ras at the membrane. This suggests a way in which a cell can use the EGFR proliferation pathway in the absence of EGF,⁴² which may have interesting ramifications in cancer.

Mannick et al. identified S-nitrosylation as a regulatory step in caspase activity.^{43,44} Nitrosylation of a thiol in the active site leads to mitochondrial localization and inactivation of caspase-3 zymogens. Fas signaling activates caspase-3 by denitrosylation of the active site thiol and translocation to the cytoplasm during the initiation of apoptosis. Nitrosylation also participates in the regulation of the apoptotic signaling through the modification of glyceraldehyde-3-phosphate dehydrogenase (GADPH). Sen et al. reported that nitric oxide produced in response to a broad range of apoptotic stimuli nitrosylates GADPH, abolishing its normal metabolic function in glycolysis and enabling it to complex with an E-3 ubiquitin ligase. The complex GADPH/E3 ubiquitin ligase translocates to the nucleus where it stimulates apoptosis.⁴⁵ In addition, Talbott et al. reported S-nitrosation as one of the mechanism of regulation of death receptor-mediated apoptosis.⁴⁶ The FLICE-inhibitory protein is a regulator on the NF- κ B/caspase-8 death pathway, and its cleavage and processing is dependent on nitrosylation in cysteine residues 254 and 259.

Oxidative modifications by peroxynitrite

Further nitric oxide mediated post-translational modifications occur though the formation of other reactive nitrogen species. Nitric oxide produces peroxynitrite through the diffusion-limited reaction with superoxide.⁴⁷ This molecule gained attention when, after studying the ability of superoxide dismutase to prevent injury in many disease states, Beckman et al identified peroxynitrite as the source of this damage.^{48,49} In an aqueous environment, the protonated peroxynitrous acid forms and can rapidly decompose into nitrogen dioxide (*NO_2) and an extremely reactive hydroxyl radical (*OH)^{24,26} with an efficiency of approximately 30%, though these values are controversial.^{48,50,51} The hydroxyl radical is reactive enough to remove electrons from nearly any biological molecule, for instance the reaction with tyrosine creates a tyrosyl radical which can then react with nitrogen dioxide to form nitrotyrosine. In addition, peroxynitrite directly reacts with CO_2 yielding nitrosoperoxycarbonate (ONOOCO_2^-), which decomposes into nitrogen dioxide and the carbonate radical (*CO_3^-) that has a similar reactivity to the hydroxyl radical.¹ The production of these carbonate and hydroxyl radicals and peroxynitrite's ability to react strongly with metal centers^{1,50,52} caused its presence to be associated with catastrophic cytotoxic circumstances (Figure 1). Peroxynitrite was seen as too reactive to participate in signaling events and its reaction with other molecules was considered to be damage rather than an oxidative modification.

The ability of peroxynitrite to modify proteins and other macromolecules has been harnessed by the phagocytes in




Figure 1 Nitric oxide reactions and protein post-translational modifications

the immune system. Both activated macrophages and neutrophils kill invading bacteria by engulfing them and releasing oxidative species in the “respiratory burst”. After phagocytosis of antigenic particles, NADPH oxidase produces large amounts of superoxide and iNOS generates nitric oxide,^{53–56} which combine to create peroxynitrite. It was found that the bacterial-killing function of activated macrophages was dependent on the presence of peroxynitrite, and that inhibiting either superoxide or nitric oxide formation inhibits this function.⁵⁵

It is important to note that while peroxynitrite is used to protect the organism from pathogens, overproduction or dysregulation of these pathways is also damaging. The same respiratory burst used by immune cells is also responsible for chronic inflammatory responses. After observing rat models of traumatic brain injury, Hall et al. noted a six-fold increase in iNOS transcript. The subsequent increase in peroxynitrite formation switched the response from protective to neuropathogenic.¹⁹ In sickle cell disease, characterized by multiple ischemia-reperfusion like events, peroxynitrite mediated damage occurs in the microvasculature epithelium when blood flow intermittently blocked by aberrant red blood cells is restored. The oxidative damage observed in reperfusion injury, and perhaps in sickle cell disease, has been recently shown to originate from the reversal of respiratory complex 1 in mitochondria causing excess superoxide production, which could then react with nitric oxide produced by eNOS and iNOS.^{57–59}

Peroxynitrite overproduction results in tissue damage characterized by oxidation of proteins leading to loss of function. Peroxynitrite affects cellular metabolism by lipid peroxidation,⁴⁹ DNA damage,⁶⁰ and mitochondrial damage.⁶¹ While peroxynitrite may have important cellular

functions, if its levels become too high, it will cause catastrophic oxidative damage to cells and tissues. For this reason peroxynitrite has been implicated in a wide variety of disease states (Table 1).

Peroxynitrite can also oxidize thiols.⁷⁵ In fact, thiols act as scavengers of peroxynitrite because thiol oxidation is the fastest direct reaction of peroxynitrite. It is important to note that thiol oxidation and nitration/nitrosylation are not mechanistically exclusive and they are competing with each other in biological relevant conditions (Figure 1).

It had been shown that peroxynitrite leads to apoptosis and although the exact mechanism was unclear at the time, it was assumed that the oxidant caused general oxidative damage by randomly modifying macromolecules eventually leading to p38 activation and cell death.⁷⁶ Since then, evidence accumulated suggesting that peroxynitrite mediated cell death was not random death due to oxidative damage, but was a tightly regulated process.^{77–79} The Akt and MLK/MAP kinase pathways were identified as directly involved in the regulation of peroxynitrite-induced apoptosis.⁸⁰ Due to the long-lasting nature of peroxynitrite modifications, peroxynitrite activation of a regulated pathway suggests a possible mechanism by which the cells could measure levels of oxidative stress. Theoretically, the accumulation of these modifications due to continuous oxidative damage could reach a threshold where the cell should commit to programmed cell death, activating p38 and leading to apoptosis. Considering this possibility creates an interesting question: If some of the peroxynitrite-induced disease states occur in a regulated pathway, could peroxynitrite function as a signaling intermediate in normal cellular processes?

Table 1 A short list of diseases and peroxynitrite's pathological role

Disease	Peroxynitrite mediated alterations	Target	Source
Amyotrophic lateral sclerosis (ALS)	Apoptosis of motor neurons	Tyrosine nitration	Zhu et al. ⁶²
Parkinson's disease	Cu,Zn-SOD damage	Tyrosine nitration	Kamisaki et al. ⁶³
	General mitochondrial damage	Iron/sulfur cluster reactions	Csermely et al. ⁶⁴
	Tyrosine hydroxylase impairment	Tyrosine nitration	Stebbins et al. ⁶⁵
Diabetes	α -synuclein modifications	Tyrosine nitration	Wiech et al. ⁶⁶
	DNA damage/binding	Guanine nitration	Franco et al. ⁶⁷
	Flow mediated dilation disruption	Tyrosine nitration	Cassuto et al. ²³
Rheumatoid arthritis	H2A histone modification, induction of antibody response	Tyrosine nitration	Franco and Estevez ⁶⁸
Lupus erythematosus	H1 histone modification, induction of antibody response	Tyrosine nitration	Franco and Estevez, ⁶⁸ Yamakura et al. ⁶⁹
Vitiligo	Mitochondrial DNA changes, immunogenic response	Guanine nitration	Jin et al. ⁷⁰
Liver diseases	Broad spectrum protein nitration, Conversion of Hsp90 to toxic species	Tyrosine nitration	Blanchard-Fillion et al. ⁷¹
Atherosclerosis	Prostaglandin synthase impairment	Tyrosine nitration	Deeb et al. ¹⁸
	LDL oxidation and accumulation	Tyrosine nitration	Przedborski et al. ⁷²
Pulmonary hypertension	Hyperactive eNOS secondary to caveolin depletion impairs PKG activity	Tyrosine nitration	Zhao et al. ²²
Chronic heart disease	Inhibition of platelet activity	Cysteine nitrosation	Tripathi et al. ⁷³
	Low serum Ferroxidase I activity	Tyrosine nitration	Khan et al. ⁷⁴

Tyrosine nitration

To help answer that question, we must look at another type of nitric oxide based, peroxynitrite mediated post-translational modification—protein nitration, one of the best characterized peroxynitrite modifications. Originally identified as a footprint of peroxynitrite presence, nitration of tyrosine is the post-translational modification seen in many of the diseases in which peroxynitrite is produced (Table 1).¹ Nitrotyrosine is present in a relatively small number of proteins, and occurs in limited positions on these proteins.¹ Recently, protein nitration has been gaining traction as a signal transduction mechanism, as suggested many years ago by Ischiropoulos and Al-Mehdi.⁸¹ We have previously shown that peroxynitrite-induced apoptosis can be prevented by tyrosine-containing peptides that scavenge the products of peroxynitrite decomposition, preventing nitration of proteins.⁸² Traditionally, nitration is seen as an irreversible process, making it poorly suited for the dynamics of signal transduction.⁸³

However, in some circumstances cells need long term signaling. Nitration is then a potential mechanism to engender lasting alterations to cell functionality, such as memory formation, cellular differentiation, or in major metabolic shifts. Growing evidence shows that post-translational modification of proteins by nitration can be used to modulate their activity in ways that other redox signals cannot. We have recently shown that nitrotyrosine levels are dynamically regulated during fetal heart development.⁸⁴ This temporal pattern does not correlate with apoptosis, but with myocyte differentiation. The permanent changes that happen in development are well suited to a signal

transduction mechanism that is not readily reversible. In 2013, researchers reported that the activity of hydroxypyruvate reductase from *Arabidopsis* plants was regulated by site-specific tyrosine nitration.⁸⁵ This loss of function modification was reported in a normal, no-stress condition, and no effect was observed after exposure to hydrogen peroxide, suggesting nitration was the intended functional modification, not a by-product of other reactive species. Finding pathways such as this where nitration is used in normal conditions supports the hypothesis that nitration can be a controlled mechanism, and not only a sign of oxidative damage. In addition, nitration of proteins in liver caused by acetaminophen is catalyzed by superoxide dismutase (SOD).^{86–88} Gene deletion of SOD prevents both nitration and damage in the liver produced by high doses of acetaminophen. These results suggest that in addition to the more accepted radical unregulated oxidative mechanisms, endogenous tyrosine nitration may be catalyzed, increasing the level of regulation.

Furthermore, Kamasaki et al. reported the ability of spleen and lung homogenates to reduce nitrotyrosine in a nitrated BSA sample.⁸⁹ They proposed the presence of a “nitrotyrosine denitrase”, although were unable to isolate the source of this activity. Recently, Deeb et al. characterized this denitrase, showing that it rapidly reversed COX-1 heme-dependent nitration in a regulated and substrate-selective manner.³ Using more modern techniques they were able to isolate, identify and quantify the denitrase in several tissue types and elucidate methods for regulation of its activity. These findings indicate that nitration is not as permanent as once believed, but a dynamic, well regulated process very suitable for signal transduction.

Nitration of heat shock protein 90

We identified heat shock protein 90 (Hsp90) as a target of post-translational nitration in PC12 cells after incubation with peroxynitrite and cultured motor neurons deprived of trophic factors using monoclonal antibodies specific only to the nitrated form of the protein.⁶⁹ Hsp90 has multiple cellular functions and is one of the most common proteins in the cell, making up 1–2% of the total protein profile in the cytoplasm. This chaperone has a very broad range of functions: protein folding, stabilization, translocation, activity modulation, and degradation.^{90–92}

Further study revealed that nitration of a single tyrosine residue on Hsp90 (Y33 or Y56) was necessary and sufficient to cause a gain of function converting this pro-survival chaperone into a pro-apoptotic protein.⁶² Nitrated Hsp90 has been identified in the spinal cords of patients suffering from sporadic ALS and in SOD mutant ALS model mice.^{62,93} Interestingly, the amount of nitrated Hsp90 also correlates with disease progression, increasing as the disease progresses, suggesting that the gain of apoptotic function by nitrated Hsp90 participates in the progressive motor neuron death characteristic of this disease (Dennys, Trias, Ramdial and Estevez, unpublished observations).

Nitration has been reported in many disease states and a large body of research has been focused on preventing or reducing these reactions, but relatively little intellectual attention has been directed towards the endogenous signaling function of these reactions. If considered in this light, this oxidative post-translational modification can be seen as something other than a stressor for the cell, but critical cellular mechanisms for maintenance and normal function. Understanding the relative proportion of the different nitration mechanisms in normal conditions, as well as how nitration is regulated could result in a better understanding of the pathogenic mechanisms activated in many disease states. It may also shed some light on the general failure of antioxidant therapies.

Author contributions: All authors participated in the writing of this manuscript.

ACKNOWLEDGEMENTS

This article was supported by NIH Grants NS36761 (to AGE).

REFERENCES

- Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 2007;**8**:315–424
- Gow AJ, Duran D, Malcolm S, Ischiropoulos H. Effects of peroxynitrite-induced protein modifications on tyrosine phosphorylation and degradation. *FEBS Lett* 1996;**385**:63–6
- Deeb RS, Nuriel T, Cheung C, Summers B, Lamon BD, Gross SS, Hajjar DP. Characterization of a cellular denitrase activity that reverses nitration of cyclooxygenase. *Am J Physiol Heart Circ Physiol* 2013;**305**:H687–98
- Smallwood HS, Lourette NM, Boschek CB, Bigelow DJ, Smith RD, Pasatolic L, Squier TC. Identification of a denitrase activity against calmodulin in activated macrophages using high-field liquid chromatography – FTICR mass spectrometry. *Biochemistry* 2007;**46**:10498–505
- Irie Y, Saeki M, Kamisaki Y, Martin E, Murad F. Histone H1.2 is a substrate for denitrase, an activity that reduces nitrotyrosine immunoreactivity in proteins. *Proc Natl Acad Sci USA* 2003;**100**:5634–9
- Shi Q, Xu H, Yu H, Zhang N, Ye Y, Estevez AG, Deng H, Gibson GE. Inactivation and reactivation of the mitochondrial alpha-ketoglutarate dehydrogenase complex. *J Biol Chem* 2011;**286**:17640–8
- Neumann H, Hazen JL, Weinstein J, Mehl RA, Chin JW. Genetically encoding protein oxidative damage. *J Am Chem Soc* 2008;**130**:4028–33
- Franco MC, Ye Y, Refakis CA, Feldman JL, Stokes AL, Basso M, Melero Fernandez de Mera RM, Sparrow NA, Calingasan NY, Kiaei M, Rhoads TW, Ma TC, Grumet M, Barnes S, Beal MF, Beckman JS, Mehl R, Estevez AG. Nitration of Hsp90 induces cell death. *Proc Natl Acad Sci USA* 2013;**110**:E1102–11
- DiDonato JA, Aulak K, Huang Y, Wagner M, Gerstenecker G, Topbas C, Gogonea V, DiDonato AJ, Tang WH, Mehl RA, Fox PL, Plow EF, Smith JD, Fisher EA, Hazen SL. Site-specific nitration of apolipoprotein A-I at tyrosine 166 is both abundant within human atherosclerotic plaque and dysfunctional. *J Biol Chem* 2014;**289**:10276–92
- Butterfield DA, Perluigi M, Reed T, Muharib T, Hughes CP, Robinson RA, Sultana R. Redox proteomics in selected neurodegenerative disorders: from its infancy to future applications. *Antioxid Redox Signal* 2012;**17**:1610–55
- Lopez-Sanchez LM, Lopez-Pedraza C, Rodriguez-Ariza A. Proteomic approaches to evaluate protein S-nitrosylation in disease. *Mass Spectrom Rev* 2014;**33**:7–20
- Mermelekas G, Makridakis M, Koeck T, Vlahou A. Redox proteomics: from residue modifications to putative biomarker identification by gel- and LC-MS-based approaches. *Expert Rev Proteomics* 2013;**10**:537–49
- Peinado JR, Diaz-Ruiz A, Fruhbeck G, Malagon MM. Mitochondria in metabolic disease: getting clues from proteomic studies. *Proteomics* 2014;**14**:452–66
- Murad F. Nobel lecture: discovery of some of the biological effects of nitric oxide and its role in cell signaling [Web] Nobelprize.org; Nobel Media, Available from: http://www.nobelprize.org/nobel_prizes/medicine/laureates/1998/murad-lecture.html%3E (1998, accessed 29 November 2014)
- Murad F. Signal transduction using nitric oxide and cyclic guanosine monophosphate. *JAMA* 1996;**279**:1189
- Jin XY, Lee SH, Kim JY, Zhao Y, Park E, Lee B, Nan J, Song K, Ko G, Sohn DH. Polyozellin inhibits nitric oxide production by down-regulating LPS-induced activity of NF-kappaB and SAPK/JNK in RAW 264.7 cells. *Planta Med* 2006;**72**:857–9
- Jung CH, Jung H, Shin YC, Park JH, Jun CY, Kim HM, Yim HS, Shin MG, Bae HS, Kim SH, Ko SG. *Eleutherococcus senticosus* extract attenuates LPS-induced iNOS expression through the inhibition of Akt and JNK pathways in murine macrophage. *J Ethnopharmacol* 2007;**113**:183–7
- Deeb RS, Shen H, Gamss C, Gavrilova T, Summers BD, Kraemer R, Hao G, Gross SS, Laine M, Maeda N, Hajjar DP, Upmacis RK. Inducible nitric oxide synthase mediates prostaglandin h2 synthase nitration and suppresses eicosanoid production. *Am J Pathol* 2006;**168**:349–62
- Hall ED, Wang JA, Miller DM. Relationship of nitric oxide synthase induction to peroxynitrite-mediated oxidative damage during the first week after experimental traumatic brain injury. *Exp Neurol* 2012;**238**:176–82
- Kumar A, Chen SH, Kadiiska MB, Hong JS, Zielonka J, Kalyanaraman B, Mason RP. Inducible nitric oxide synthase is key to peroxynitrite-mediated, LPS-induced protein radical formation in murine microglial BV2 cells. *Free Radic Biol Med* 2014;**73**:51–9
- Shelton JL, Wang L, Cepinskas G, Sandig M, Scott JA, North ML, Inculet R, Mehta S. Inducible NO synthase (iNOS) in human neutrophils but not pulmonary microvascular endothelial cells (PMVEC) mediates septic protein leak in vitro. *Microvasc Res* 2007;**74**:23–31
- Zhao Y-Y, Zhao YD, Mirza MK, Huang JH, Potula H-HSK, Vogel SM, Brovkovich V, Yuan JXJ, Wharton J, Malik AB. Persistent eNOS activation secondary to caveolin-1 deficiency induces pulmonary hypertension in mice and humans through PKG nitration. *J Clin Invest* 2009;**119**:2009
- Cassuto J, Dou H, Czika I, Szabo A, Patel VS, Kamath V, de Chantemele EB, Feher A, Romero MJ, Bagi Z. Peroxynitrite disrupts

- endothelial caveolae leading to eNOS uncoupling and diminished flow-mediated dilation in coronary arterioles of diabetic patients. *Diabetes* 2014;**63**:1381+
24. Francis SH, Busch JL, Corbin JD, Sibley D. cGMP-dependent protein kinases and cGMP phosphodiesterases in nitric oxide and cGMP action. *Pharmacol Rev* 2010;**62**:525–63
 25. Estevez AG, Spear N, Thompson JA, Cornwell TL, Radi R, Barbeito L, Beckman JS. Nitric oxide-dependent production of cGMP supports the survival of rat embryonic motor neurons cultured with brain-derived neurotrophic factor. *J Neurosci* 1998;**18**:3708–14
 26. McDonald LJ, Murad F. Nitric oxide and cGMP signaling. *Adv Pharmacol* 1995;**34**:263–75
 27. Chiueh CC. Neuroprotective properties of nitric oxide. *Ann N Y Acad Sci* 1999;**890**:301–11
 28. Denninger JW, Marletta MA. Review: guanylate cyclase and the .NO/ cGMP signaling pathway. *BBA - Bioenergetics* 1999;**1411**:334–50
 29. Garthwaite J. Concepts of neural nitric oxide-mediated transmission. *Eur J Neurosci* 2008;**27**:2783–802
 30. Krumenacker JS, Hanafy KA, Murad F. Regulation of nitric oxide and soluble guanylyl cyclase. *Brain Res Bull* 2004;**62**:505–15
 31. Martin E, Berk A, Tsai AL, Murad F. Soluble guanylyl cyclase: the nitric oxide receptor. *Methods Enzymol* 2005;**396**:478–92
 32. Reid MB. Role of nitric oxide in skeletal muscle: synthesis, distribution and functional importance. *Acta Physiol Scand* 1998;**162**:401–9
 33. Stamler JS, Simon DI, Osborne JA, Mullins ME, Jaraki O, Michel T, Singel DJ, Loscalzo J. S-Nitrosylation of Proteins with Nitric Oxide: Synthesis and Characterization of Biologically Active Compounds. *Proc Natl Acad Sci USA* 1992;**89**:444
 34. Stamler JS, Lamas S, Fang FC. Nitrosylation: the prototypic redox-based signaling mechanism. *Cell* 2001;**106**:675
 35. Doulias P-T, Greene JL, Greco TM, Tenopoulou M, Seeholzer SH, Dunbrack RL, Ischiropoulos H. Structural profiling of endogenous S-nitrosocysteine residues reveals unique features that accommodate diverse mechanisms for protein S-nitrosylation. *Proc Natl Acad Sci USA* 2010;**107**:16958
 36. Heinrich TA, Silva RS, Miranda KM, Switzer CH, Wink DA, Fukuto JM. Biological nitric oxide signalling: chemistry and terminology. *Br J Pharmacol* 2013;**169**:1417
 37. Evangelista AM, Murphy E, Kohr MJ. S-Nitrosylation: specificity, occupancy, and interaction with other post-translational modifications. *Antioxid Redox Signal* 2013;**19**:1209
 38. Martinez-Ruiz A, Araujo IM, Izquierdo-Alvarez A, Hernansanz-Agustin P, Lamas S, Serrador JM. Specificity in S-nitrosylation: a short-range mechanism for no signaling? *Antioxid Redox Signal* 2013;**19**:1220–35
 39. Yun H-Y, Gonzalez-Zulueta M, Dawson VL, Dawson TM. Nitric oxide mediates N-methyl-D-aspartate receptor-induced activation of p21 ras. *Proc Natl Acad Sci USA* 1998;**95**:5773
 40. Lipton SA, Choi Y-B, Pan Z-H, Lei SZ, Chen H-SV, Sucher NJ, Loscalzo J, Singel DJ, Stamler JS. A redox-based mechanism for the neuroprotective and neurodestructive effects of nitric oxide and related nitroso-compounds. *Nature* 1993;**364**:626–32
 41. Lander HM, Ogiste JS, Teng KK, Novogrodsky A. p21ras as a common signaling target of reactive free radicals and cellular redox stress. *J Biol Chem* 1995;**270**:21195–8
 42. Batista WL, Ogata FT, Curcio MF, Miguel RB, Arai RJ, Matsuo AL, Moraes MS, Stern A, Monteiro HP. S-nitrosoglutathione and endothelial nitric oxide synthase-derived nitric oxide regulate compartmentalized ras S-nitrosylation and stimulate cell proliferation. *Antioxid Redox Signal* 2013;**18**:221–38
 43. Mannick JB. Fas-induced caspase denitrosylation. *Science* 1999;**284**:651–4
 44. Mannick JB, Schonhoff C, Papeta N, Ghafourifar P, Szibor M, Fang K, Gaston B. S-Nitrosylation of mitochondrial caspases. *J Cell Biol* 2001;**154**:1111–6
 45. Sen N, Hara MR, Kornberg MD, Cascio MB, Bae BI, Shahani N, Thomas B, Dawson TM, Dawson VL, Snyder SH, Sawa A. Nitric oxide-induced nuclear GAPDH activates p300/CBP and mediates apoptosis. *Nat Cell Biol* 2008;**10**:866–73
 46. Talbott SJ, Luanpitpong S, Stehlik C, Azad N, Iyer AK, Wang L, Rojanasakul Y. S-nitrosylation of FLICE inhibitory protein determines its interaction with RIP1 and activation of NF-kappaB. *Cell Cycle* 2014;**13**:1948–57
 47. Nauser T, Koppenol WH. The rate constant of the reaction of superoxide with nitrogen monoxide: approaching the diffusion limit. *J Phys Chem A* 2002;**106**:4084
 48. Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent Hydroxyl Radical Production by Peroxynitrite: Implications for Endothelial Injury from Nitric Oxide and Superoxide. *Proc Natl Acad Sci USA* 1990;**87**:1620
 49. Radi R, Beckman JS, Bush KM, Freeman BA. Peroxynitrite-induced membrane lipid peroxidation: the cytotoxic potential of superoxide and nitric oxide. *Arch Biochem Biophys* 1991;**288**:481–7
 50. Forman HJ, Ursini F, Maiorino M. An overview of mechanisms of redox signaling. *J Mol Cell Cardiol* 2014;**73**:2–9
 51. Koppenol WH, Bounds PL, Nauser T, Kissner R, Ruegger H. Peroxynitrous acid: controversy and consensus surrounding an enigmatic oxidant. *Dalton Trans* 2012;**41**:13779–87
 52. Beckman JS, Carson M, Smith CD, Koppenol WH. ALS, SOD and peroxynitrite. *Nature* 1993;**364**:584
 53. Gilbert RS, Herschman HR. Transforming growth factor beta differentially modulates the inducible nitric oxide synthase gene in distinct cell types. *Biochem Biophys Res Commun* 1993;**195**:380–4
 54. Carreras MC, Pargament GA, Catz SD, Poderoso JJ, Boveris A. Kinetics of nitric oxide and hydrogen peroxide production and formation of peroxynitrite during the respiratory burst of human neutrophils. *FEBS Lett* 1994;**341**:65–8
 55. Allen RG, Lafuse WP, Powell ND, Webster Marketon JJ, Stiner-Jones LM, Sheridan JF, Bailey MT. Stressor-induced increase in microbicidal activity of splenic macrophages is dependent upon peroxynitrite production. *Infect Immun* 2012;**80**:3429–37
 56. Dahlgren C, Karlsson A. Respiratory burst in human neutrophils. *J Immunol Methods* 1999;**232**:3–14
 57. Chouchani ET, Pell VR, Gaude E, Aksentijevic D, Sundier SY, Robb EL, Logan A, Nadtochiy SM, Ord EN, Smith AC, Eyassu F, Shirley R, Hu CH, Dare AJ, James AM, Rogatti S, Hartley RC, Eaton S, Costa AS, Brookes PS, Davidson SM, Duchon MR, Saeb-Parsy K, Shattock MJ, Robinson AJ, Work LM, Frezza C, Krieg T, Murphy MP. Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 2014;**515**:431–5
 58. Bank N, Kiroychewa M, Ahmed F, Anthony GM, Fabry ME, Nagel RL, Singhal PC. Peroxynitrite formation and apoptosis in transgenic sickle cell mouse kidneys. *Kidney Int* 1998;**54**:1520–8
 59. Chirico EN, Pialoux V. Role of oxidative stress in the pathogenesis of sickle cell disease. *IUBMB Life* 2012;**64**:72–80
 60. Groves JT, Marla SS. Peroxynitrite-induced DNA strand scission mediated by a manganese porphyrin. *J Am Chem Soc* 1995;**117**:9578–9
 61. Radi R, Cassina A, Hodara R, Quijano C, Castro L. Peroxynitrite reactions and formation in mitochondria. *Free Radic Biol Med* 2002;**33**:1451–64
 62. Zhu JH, Zhang XM, McClung JP, Lei XG. Impact of Cu,Zn-superoxide dismutase and Se-dependent glutathione peroxidase-1 knockouts on acetaminophen-induced cell death and related signaling in murine liver. *Exp Biol Med* 2006;**231**:1726–32
 63. Kamisaki Y, Wada K, Bian K, Balabanli B, Davis K, Martin E, Behbod F, Lee Y-C, Murad F. An activity in rat tissues that modifies nitrotyrosine-containing proteins. *Proc Natl Acad Sci USA* 1998;**95**:11584
 64. Csermely P, Schnaider T, Soti C, Prohászka Z, Nardai G. The 90-kDa molecular chaperone family: structure, function, and clinical applications. A comprehensive review. *Pharmacol Therapeut* 1998;**79**:129–68
 65. Stebbins CE, Russo AA, Schneider C, Rosen N, Hartl FU, Pavletich NP. Crystal structure of an Hsp-90-geldanamycin complex: targeting of a protein chaperone by an antitumor agent. *Cell* 1997;**89**:239–50
 66. Wiech H, Buchner J, Zimmermann R, Jakob U. Hsp90 chaperones protein folding in vitro. *Nature* 1992;**358**:169–70
 67. Franco MC, Ye Y, Refakis CA, Feldman JL, Stokes AL, Basso M, Melero Fernandez de Mera RM, Sparrow NA, Calingasan NY, Kiaei M, Rhoads TW, Ma TC, Grumet M, Barnes S, Beal MF, Beckman JS, Mehl R,

- Estevez AG. Nitration of Hsp90 induces cell death. *Proc Natl Acad Sci USA* 2013;**110**:E1102–11
68. Franco MC, Estevez AG. Tyrosine nitration as mediator of cell death. *Cell Mol Life Sci* 2014;**71**:3939–50
 69. Yamakura F, Matsumoto T, Ikeda K, Taka H, Fujimura T, Murayama K, Watanabe E, Tamaki M, Imai T, Takamori K. Nitrated and oxidized products of a single tryptophan residue in human Cu,Zn-superoxide dismutase treated with either peroxynitrite-carbon dioxide or myeloperoxidase-hydrogen peroxide-nitrite. *J Biochem* 2005;**138**:57–69
 70. Jin H, Kanthasamy A, Ghosh A, Anantharam V, Kalyanaraman B, Kanthasamy AG. Mitochondria-targeted antioxidants for treatment of Parkinson's disease: preclinical and clinical outcomes. *Biochim Biophys Acta* 2014;**1842**:1282–94
 71. Blanchard-Fillion B, Souza JM, Friel T, Jiang GCT, Vrana K, Sharov V, Barrón L, Schöneich C, Quijano C, Alvarez B, Radi R, Przedborski S, Fernando GS, Horwitz J, Ischiropoulos H. Nitration and inactivation of tyrosine hydroxylase by peroxynitrite. *J Biol Chem* 2001;**276**:46017–23
 72. Przedborski S, Chen Q, Vila M, Giasson BI, Djaldatti R, Vukosavic S, Souza JM, Jackson-Lewis V, Lee VMY, Ischiropoulos H. Oxidative post-translational modifications of α -synuclein in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) mouse model of Parkinson's disease. *J Neurochem* 2001;**76**:637–40
 73. Tripathi P, Moinuddin, Dixit K, Mir AR, Habib S, Alam K, Ali A. Peroxynitrite modified DNA presents better epitopes for anti-DNA autoantibodies in diabetes type 1 patients. *Cell Immunol* 2014;**290**:30–8
 74. Khan MA, Dixit K, Uddin M, Malik A, Alam K. Role of peroxynitrite-modified H2A histone in the induction and progression of rheumatoid arthritis. *Scand J Rheumatol* 2012;**41**:426–33
 75. Ahmad R, Ahsan H. Role of peroxynitrite-modified biomolecules in the etiopathogenesis of systemic lupus erythematosus. *Clin Exp Med* 2014;**14**:1–11
 76. Al-Shobaili HA, Rasheed Z. Mitochondrial DNA acquires immunogenicity on exposure to nitrosative stress in patients with vitiligo. *Hum Immunol* 2014;**75**:1053–61
 77. Abdelmegeed MA, Song BJ. Functional roles of protein nitration in acute and chronic liver diseases. *Oxid Med Cell Longev* 2014;**2014**:149627
 78. Leeuwenburgh C, Hardy MM, Hazen SL, Wagner P, Oh-ishi S, Steinbrecher UP, Heinecke JW. Reactive nitrogen intermediates promote low density lipoprotein oxidation in human atherosclerotic intima. *J Biol Chem* 1997;**272**:1433–6
 79. Olas B, Nowak P, Wachowicz B. Resveratrol protects against peroxynitrite-induced thiol oxidation in blood platelets. *Cell Mol Biol Lett* 2004;**9**:577–87
 80. Quijano C, Alvarez B, Gatti RM, Augusto O, Radi R. Pathways of peroxynitrite oxidation of thiol groups. *Biochem J* 1997;**322**:167–73
 81. Cabassi A, Binno SM, Tedeschi S, Ruzicka V, Dancelli S, Rocco R, Vicini V, Coghi P, Regolisti G, Montanari A, Fiaccadori E, Govoni P, Piepoli M, de Champlain J. Low serum ferroxidase I activity is associated with mortality in heart failure and related to both peroxynitrite-induced cysteine oxidation and tyrosine nitration of ceruloplasmin. *Circ Res* 2014;**114**:1723–32
 82. Jope RS, Zhang L, Song L. Peroxynitrite modulates the activation of p38 and extracellular regulated kinases in PC12 cells. *Arch Biochem Biophys* 2000;**376**:365–70
 83. Estévez AG, Radi R, Barbeito L, Shin JT, Thompson JA, Beckman JS. Peroxynitrite-Induced Cytotoxicity in PC12 Cells: evidence for an apoptotic mechanism differentially modulated by neurotrophic factors. *J Neurochem* 1995;**65**:1543–50
 84. Raoul C, Estevez AG, Nishimune H, Cleveland DW, deLapeyriere O, Henderson CE, Haase G, Pettmann B. Motoneuron death triggered by a specific pathway downstream of Fas: potentiation by ALS-linked SOD1 mutations. *Neuron* 2002;**35**:1067–83
 85. Spear N, Estévez AG, Barbeito L, Beckman JS, Johnson GVW. Nerve growth factor protects PC12 cells against peroxynitrite-induced apoptosis via a mechanism dependent on phosphatidylinositol 3-kinase. *J Neurochem* 1997;**69**:53–9
 86. Shacka JJ, Sahawneh MA, Gonzalez JD, Ye YZ, D'Alessandro TL, Estevez AG. Two distinct signaling pathways regulate peroxynitrite-induced apoptosis in PC12 cells. *Cell Death Differ* 2006;**13**:1506–14
 87. Ischiropoulos H, Al-Mehdi AB. Peroxynitrite-mediated oxidative protein modifications. *FEBS Lett* 1995;**364**:279–82
 88. Ye Y, Quijano C, Robinson KM, Ricart KC, Strayer AL, Sahawneh MA, Shacka JJ, Kirk M, Barnes S, Accavitti-Loper MA, Radi R, Beckman JS, Estevez AG. Prevention of peroxynitrite-induced apoptosis of motor neurons and PC12 cells by tyrosine-containing peptides. *J Biol Chem* 2007;**282**:6324–37
 89. Janssen-Heininger YMW, Mossman BT, Heintz NH, Forman HJ, Kalyanaraman B, Finkel T, Stamler JS, Rhee SG, van der Vliet A. Review article: redox-based regulation of signal transduction: principles, pitfalls, and promises. *Free Radic Biol Med* 2008;**45**:1–17
 90. Viera L, Radmilovich M, Vargas MR, Dennys CN, Wilson L, Barnes S, Franco MC, Beckman JS, Estevez AG. Temporal patterns of tyrosine nitration in embryo heart development. *Free Radic Biol Med* 2013;**55**:101–8
 91. Corpas FJ, Leterrier M, Begara-Morales JC, Valderrama R, Chaki M, Lopez-Jaramillo J, Luque F, Palma JM, Padilla MN, Sanchez-Calvo B, Mata-Perez C, Barroso JB. Inhibition of peroxisomal hydroxypyruvate reductase (HPR1) by tyrosine nitration. *Biochim Biophys Acta* 2013;**1830**:4981–9
 92. Lei XG, Zhu J-H, McClung JP, Aregullin M, Roneker CA. Mice deficient in Cu,Zn-superoxide dismutase are resistant to acetaminophen toxicity. *Biochem J* 2006;**399**:455–61
 93. Zhu JH, Zhang X, Roneker CA, McClung JP, Zhang S, Thannhauser TW, Ripoll DR, Sun Q, Lei XG. Role of copper,zinc-superoxide dismutase in catalyzing nitrotyrosine formation in murine liver. *Free Radic Biol Med* 2008;**45**:611–8